

"A process for preparing cephalosporin derivatives"

5 The invention relates to a process for the preparation of 3-
cephalosporin C derivatives which are used in the preparation of β -
lactam antibiotics. In particular, the invention relates to a process for
the preparation of 3-thiolated derivatives of 3-acetoxy-methyl-7-amino-
ceph-3-em-carboxylic acid.

Background

10

All cephalosporins used in therapeutic applications are semi-synthetic
and are produced by modifying the basic β -lactam structure in the
material obtained from a fermentation broth of *Acremonium*
chrysogenum or, after chemical transformation, from the products
15 obtained from fermentation broths of *Penicillium chrysogenum* using
different precursors.

Typically, cephalosporin C [3-acetoxymethyl-7-(D-5-amino-5-carboxy
pentan amido)-ceph-3-em-4-carboxylic acid] is converted into 3-
20 acetoxymethyl-7-amino-ceph-3-em-carboxylic acid, usually known as
7-amino cephalosporanic acid (7-ACA), by removing the lateral
aminoadipic chain of the β -lactam ring. The 7-ACA is purified and
crystallised, and is then used as starting material for subsequent
modifications at the 7- and 3- position. 7-ACA is the base building
25 block used in the synthesis of many important semi-synthetic
cephalosporin antibiotics that are of current interest in the
biopharmaceutical industry.

3-thiomethyl Cephalosporin C derivatives have been disclosed in US 3,278,531; US 3,516,997; US 3,647,788; GB 1,400,804; GB 1,565,053 and GB 1,566,515. A few examples of 3'-thiomethyl glutaryl 7-ACA derivatives are described in WO-A-9535020 and EP 0846695.

5

7-ACA is the most widely used intermediate for reaction with heterocyclic thiols as it can be obtained either chemically or enzymatically on an industrial scale. This has been disclosed in several patents including US 3,502,665; US 3,954,745; US 3,516,997; 10 US 3,979,383; US 4,115,645; US 4,317,907; US 5,387,679; JP 55,139,327; EP 0167651 and WO-A-9302085.

US 5,387,679 describes the reaction of 7-amino cephalosporanic acid with 2-mercapto-5-methyl-1,3,4-thiadiazole (MMTD) in the presence of 15 sodium bicarbonate in aqueous acetone, at pH 6-7. A yield of about 60-65 % is achieved.

In US 4,317,907, the yield of the above reaction in anhydrous medium was increased to about 86 % when acetic acid or nitromethanol was 20 used as a solvent, in the presence of boron trifluoride or boron trifluoride etherate. However the purity is low, 80 % max, as it contains unreacted 7-ACA and degradation products. In WO-A-9302085 the use of dialkyl carbonate in the presence of a dialkyl carbonate trifluoride complex and an aliphatic acid increased the yield 25 to a maximum of 89%. However, WO-A-9302085 shows the problems associated with using toxic and very expensive gas such as BF_3 and the handling of waste effluent which contains borides and fluoroborides.

The deacylation of cephalosporin C, i.e., the elimination of the 7'-lateral side chain, is usually carried out chemically, for example using nitrosyl chloride in formic acid in the presence of acetonitrile (US 3,367,933). Another method of deacylation involves the protection of the carboxyl group of aminopentanoic chain, reaction with phosphorus pentachloride at -55°C and subsequent hydrolysis at low temperature with a mixture of water and methanol (BE 718,824).

These known methods must generally be carried out at low temperatures, and require the use of costly and toxic solvents or reagents, consequently they may have a serious environmental impact. In addition, because of the chemical instability of the β -lactam nucleus and the reactivity of the groups present in positions 3 and 7 of the ring, special reaction conditions must be applied, which makes the process on an industrial scale complex.

To overcome the drawbacks of the chemical route to 7-ACA, alternative enzymatic cleavage of cephalosporin C has been described. Direct one-step removal of the lateral 7-aminoadipic side-chain of cephalosporin C is possible by using specific cephalosporin acylases (FR 2,241,557; US 4,774,179; EP 283,248; WO-A-9512680; WO-A-9616174). These processes, however, are often not reproducible and are characterised by low yields and lengthy reaction times as described in US 5,296,358. No industrial application of this technology (single-step conversion of cephalosporin C to 7-ACA) has been reported at this time (Parmar et al, Crit. Rev. Biotechnol. 18, 1, 1998).

On the other hand, processes that transform the cephalosporin C into 7-ACA by means of two enzymatic steps are important from an

industrial point of view. The first stage consists of using a D-amino acid oxidase (E.C. 1.4.3.3, hereinbelow indicated as DAAO) from different sources (*Trigonopsis variabilis*, GB 1,272,769; *Rhodotorula gracilis*, EP 0,517,200; or *Fusarium solari* M-0718, EP 0,364,275).

5 DAAO oxidises the lateral D-5-amido-carboxypentanoyl chain of cephalosporin C in the presence of molecular oxygen, to produce 7 β -(5-carboxy-5-oxopent-amido)-ceph-3-em-carboxylic acid (or α -ketoadipyl-7-aminocephalo-sporanic acid, hereinbelow indicated as α -ketoadipyl-7-ACA) and hydrogen peroxide, which chemically
10 decarboxylate the α -ketoadipyl-7-ACA to 7 β -(4-carboxy butanamido)-ceph-3-em-4-carboxylic acid (or glutaryl-7-aminocephalosporanic acid, hereinbelow indicated as GL-7-ACA).

In a second stage, a specific acylase for GL-7-ACA, glutaryl-7-ACA
15 acylase (E.C. 3.5.1.3), is used, for example that of a *Pseudomonas* type microorganism (US 3,960,662, EP 0496993) over expressed in *E. coli*, which deacylates the GL-7-ACA into 7-ACA.

This two-step enzymatic process for obtaining 7-ACA has been used on
20 an industrial scale (Conlon et al. Biotechnol. Bioeng. 46, 510, 1995).

An environmental-friendly alternative to produce 3'-heterocyclic thiomethyl cephalosporanic acid derivatives was disclosed in EP 0846695. Chemical nucleophilic displacement of the 3-position of
25 glutaryl-7-ACA in aqueous medium is followed by an enzymatic transformation of the 3'-glutaryl-7-ACA-derivative by using the enzyme glutaryl-7-ACA acylase. The quantity of the derivative is about 65 % with no environmental impact. This procedure can be defined as an enzymatic-chemical-enzymatic (ECE) process, since the isolated GL-7-

ACA comes from a bioconversion of solubilised cephalosporin C, then GL-7-ACA is reacted with the heterocyclic thiols and finally the 3-heterocyclic thio-derivative is enzymated with GL-7-ACA acylase. The problem with this method is the need to isolate GL-7-ACA, which given its high water solubility, is technically difficult and expensive, as described in WO-A-9535020.

An additional problem is that the enzyme is only reusable for a few cycles. In fact no more than three cycles are described with 5-mercaptotriazol and MMTD, where the residual amount of MMTD after crystallisation is 2.8 mg/ml. The other two thiols used as examples are only used in one cycle due to the high amounts of the thiols remaining in solution. This is the case with 5-mercapto-1-methyltetrazole (MMTZ) or 2,5-dihydro-3-mercapto-2-methyl-5,6-dioxo-1,2,4-triazine (TTZ), which are very soluble in water and not removable by decreasing the pH. This poisoning effect of the thiol has also been described for MMTD on Penicillin G Amidase from *E. coli* CFC-04017 (Won et al, App. Biochem. Biotech. 69, 1, 1998) during the enzymatic synthesis of cefazolin. To overcome this inhibitory effect the molar ratio of MMTD/7-ACA is decreased to 1.2:1 in order to extend the lifetime of the enzyme. This low molar ratio reduces the yield of the 3-thio derivative and avoids the use of the other published process on the chemo-enzymatic synthesis of 3-modified cephalosporins, in which an enzymatic-enzymatic-chemical process (EEC) is proposed, using D-amino acid oxidase – glutaryl-7-ACA acylase and a chemical reaction with heterocyclic thiols (Jistiz et al., J. Org. Chem. 62, 9099, 1997).

There is therefore a need for an improved process for preparing 3-cephalosporin C derivatives.


Statements of Invention

5

According to the invention there is provided an enzymatic process for preparing 3-thiolated 7-aminocephalosporanic acid derivatives comprising the steps:-


10

enzymatically converting a 3-thiolated cephalosporin C of the formula I:



15


to form a 3-thiolated -glutaryl-7-ACA of the formula II



20

and enzymatically converting a compound of formula II to form a 3-thiolated-7-ACA of the formula III

25



30

5 wherein R is a heterocyclic group comprising at least one nitrogen atom and R₁ and R₂ are both hydrogen atoms or one of them is a hydrogen atom and the other is an acyl donor.

Preferably the 3-thiolated cephalosporin C of formula I is converted into a 3-thiolated-glutaryl-7-ACA of the formula II by:-

10 reacting a compound of formula I with immobilised D-Amino acid oxidase in the presence of molecular oxygen;

separating the supported enzyme from the aqueous reaction mixture; and

15 adding hydrogen peroxide to convert the 3-thiolated- α -ketoadipyl cephalosporanic acid into a compound of formula II.

20 In one embodiment of the invention the compound of formula I is reacted with immobilised D-Amino acid oxidase at a pressure of about 2 bar absolute, a pH of from 6.0 to 8.0, and a temperature of from 20°C to 30°C for a period of from 0.5 to 3 hours.

25 Preferably the process includes the step of washing the supported enzyme with a concentrated salt solution and adding hydrogen peroxide preferably in an amount equivalent to 30 to 50 ppm to the solution thus formed.

Most preferably the process comprises the step of eliminating excess hydrogen peroxide from the solution, preferably by adding a catalyst to the solution.

5 In one embodiment of the invention the excess hydrogen peroxide is removed by adding catalase to the solution. Preferably a compound of formula II is converted into a compound of formula III by contacting a compound of formula II with immobilised glutaryl-7-ACA acylase. Most preferably the reaction to form a compound of formula III from a
10 compound of formula II is carried out at ambient pressure, at a pH of from 6.0 to 8.5 and at a temperature of from 20°C to 35°C, for a period of from 0.5 to 3 hours under an inert atmosphere.

15 In one embodiment of the invention the compound of formula III is precipitated by acidifying the reaction medium and the precipitate thus formed is subsequently washed and dried.

In another embodiment of the invention the enzymes are immobilised using a suitable cross-linker agent in a suitable solid support.
20 Preferably the enzymes are in the form of crystals of a size suitable for use as a biocatalyst. Most preferably the enzymatic processes are carried out while maintaining the enzyme in dispersion in an aqueous substrate solution.

25 In one embodiment of the invention the or each enzymatic process is carried out in a column.

Preferably the process includes the step of recovering the enzyme for reuse.

In one embodiment of the invention crystallisation of a compound of formula III is carried out at an acidic pH.

5 One aspect of the invention provides a process of the invention wherein the enzymatic conversion of a 3 thiolated cephalosporin C of the formula I to form a 3 thiolated-7-ACA of the formula III is carried out in one step.

10 In one embodiment of the invention compounds of formula I are in a solid form or in the form of a non-toxic salt thereof. Non-toxic salts may include cations used in the crystallisation of cephalosporin C which are non-toxic for humans such as zinc salts.

15 The invention also provides a process for the preparation of Cephalosporin C antibiotics and derivatives thereof comprising forming a compound of formula III and subsequent enzymation. Preferably the antibiotic is selected from any one or more of cefazolin, cefazedone, cefoperazone, cefamandol, cefatriazine, cefotiam or
20 ceftriaxone.

In one embodiment of the invention the enzymatic process for preparing 3-thiolated 7-aminocephalosporanic acid derivatives comprises the steps:-

25

reacting Cephalosporin C with a thiol compound of the general
Formula IV

R-SH

(IV)

wherein R is a heterocyclic group comprising at least one nitrogen atom,

5 to form a compound of formula I

and, after formation of the compound of formula I removing excess thiol of Formula IV.

10 Preferably the excess thiol is removed by adsorption on an anion exchange resin. Most preferably the anion exchange resin is a microporous resin having a cross-linked acrylic copolymer structure. Preferably the anion exchange resin comprises an 8% cross-linking containing functional thialkyl benzyl ammonium group. The resin may
15 be in the chloride, hydroxy, phosphate or acetate cycle.

In one embodiment of the invention the excess thiol is removed by crystallisation. Preferably crystallisation is carried out at an acidic pH. Most preferably the excess thiol is removed by crystallisation followed
20 by adsorption on an anion exchange resin.

In one embodiment of the invention the cephalosporin C is in an aqueous medium.


25 In another embodiment of the invention the cephalosporin C is in the form of a concentrated cephalosporin C solution obtained from solid cephalosporin C or from a direct or purified cephalosporin C broth. The cephalosporin C may be in the form of a concentrated cephalosporin C solution or a concentrated cephalosporin C broth.

In one embodiment of the invention the reaction process is carried out at a pH of between 5.5 and 8.0, at a temperature of from 60°C to 80°C, for a period of from 1 to 12 hours. Preferably the reaction is carried out at a pH of approximately 6.0 and at a temperature of approximately 65°C. Most preferably the thiol compound is present in an amount of between 1 and 5 mol/mol of cephalosporin C.

R may be a heterocyclic group comprising at least one nitrogen atom and optionally a sulphur or oxygen atom. Preferably R is a heterocyclic group selected from any one or more of the group comprising thienyl, diazolyl, tetrazolyl, thiazolyl, triazinyl, oxazolyl, oxadiazolyl, pyridyl, pirimidinyl, benzo thiazolyl, benzimidazolyl, benzoxazolyl, or any derivative thereof, preferably 5-methyl-1,3,4-thiadiazol-2-yl, 1-methyl-1H-tetrazol-5-yl or 1,2,5,6-tetrahydro-2-methyl-5,6-dioxo-1,2,4-triazin-3-yl.


In one embodiment of the invention compounds of formula I are in a solid form or in the form of a non-toxic salt thereof.

The invention provides a compound of the formula:-




wherein R is a heterocyclic group comprising at least one nitrogen atom, obtained by a process of the invention.

The invention also provides a compound of the formula:-



The invention provides a compound of the formula:-



wherein in formula I, R is 1,2,5,6-tetrahydro-2-methyl-5,6-dioxo-1,2,4-triazin-3-yl.

The invention also provides a process for the preparation of cephalosporin C antibiotics and derivatives thereof comprising forming a compound of formula I as hereinbefore defined and subsequent enzymation of the compound of formula I. Preferably the antibiotic is selected from any one or more of cefazolin, cefazedone, cefoperazone, cefamandol, cefatriazine, cefotiam and ceftriaxone.


Detailed description

We have found that a chemical-enzymatic-enzymatic process (CEE), in which cephalosporin C in solution is first reacted with a heterocyclic


thiol before enzymatic removal of the 7-position side chain, provides an improved process for preparing 3-cephalosporin C derivatives.

The process of the invention can be represented as follows.


First stage: Immobilised D-amino acid oxidase in the presence of O_2



3-thiolated cephalosporin C (T'X'C)




3-thiolated- α -ketoadipyl-7-ACA (T'X'K)




3-thiolated-glutaryl-7-ACA (T'X'G)

Second stage: Immobilised glutaryl-7-ACA-acylase

5



10



15

20

25

where R is a heterocyclic comprising at least a nitrogen atom with or without a sulphur or oxygen, such as thienyl, diazolyl, triazinyl, triazolyl, tetrazolyl, thiazolyl, thiadiazolyl, thiatriazolyl, oxazolyl, oxadiazolyl, pyridyl, pyrimidinyl, benzothiazolyl, benzimidazolyl, benzoxazolyl or their substituted derivatives in any of the possible positions of the above mentioned heterocyclic structure, available for substitution.

The enzymatic conversion of the compound I into the 3-thiolated glutaryl-7-ACA (compound II) is carried out in aqueous solution with a pH of about 6.5 to 8.0, preferably at pH 7.0. This avoids the problem of

increasing instability of cephalosporanic compounds at pH levels above 8. The reaction temperature can be operated from 15°C to 35°C, and is normally fixed to 20°C. The concentration of 3-thiolated cephalosporin C derivatives can vary from 35-150 mM.

5

Molecular oxygen acts as a co-substrate for the oxidative deamination. This is sparged into the reaction solution through a bottom diffuser at a flow rate from 0.01 to 1 vol/vol of solution/minute, preferably at 0.1 vvm under a suitable mechanical stirring. It will be apparent to a person skilled in the art that the rate of stirring may vary according to a wide range of design parameters and reaction characteristics. Ordinarily, the stirring would be in the range 20 – 500 rpm. This design of bioreactor is preferred to a percolation column containing the immobilised crystalline enzyme, in which it is difficult to obtain adequate transfer of oxygen, thus reducing the final yield of 3-thiolated glutaryl-7-ACA.

10
15

The time required for complete transformation is of the order of 0.5 to 3 hours, depending on the operating conditions, but is typically approximately 1 hour.

20

The conversion from compound of formula I to compound of formula II is high, and normally about 98 % with a product yield of 95 %. At the end of the reaction, the reaction solution still contains some undegraded intermediate compounds of formula IV, which are converted to compounds of formula II using external hydrogen peroxide. In addition, some compounds of formula II can remain bound to the resin in which the enzymes are immobilised.

25

To reduce the impact of these potential losses, the reactor solution is discharged with overpressure to a holding tank and the enzyme is washed with a 100 mM phosphate buffer at a pH of 7.0. Other buffer solutions with similar characteristics may be used. This buffer wash is
5 combined with the reaction solution, which mixture is then titrated with hydrogen peroxide, final concentration in solution 30-50 ppm for 30-50 minutes at about 25°C, preferably to 35 ppm for 30 minutes.

The excess of H_2O_2 is eliminated before passing to the next step by a
10 suitable enzyme, such as catalase, a suitable reducing agent such as pyruvic acid, alkaline sulphites or any other suitable catalyst in soluble or immobilised form. The enzymatic removal of H_2O_2 is preferred in this invention to preserve the quality and purity of the final product.

15 By the above treatments, the final compound (II) yield was 97-98 % which is 3 % higher than that described in EP 0496993 for the conversion of unmodified cephalosporin C into glutaryl-7-ACA.

The final solution after H_2O_2 titration is adjusted to a pH of
20 approximately 7.5 to 8.5, preferably to pH 8.0 with a concentrated organic or inorganic base, such as ammonia, before being transferred to the second bioreactor charged with a suitable wet immobilised or crystalline glutaryl-7-ACA acylase preparation. Commercially
25 available preparations, such as are available from Roche Molecular Biochemicals are suitable for the purpose. The pH is maintained by dosing the reaction with the same organic or inorganic base as above described by using an autotitrator fixed at about pH 8.0. The operating temperature may be from about 15°C to 25°C, and is preferably 20°C. The conversion time varies from 0.5 to 2 hours, depending on the

operating conditions, but usually is about 1 hour. In order to avoid microbial contamination of the enzyme preparation, a stream of nitrogen is flowed through a bottom diffuser at about 0.01 vol/vol/min under a gentle agitation.

5

After the reaction is completed, determined by the conversion of not less than 97 % of 3-thiolated glutaryl-7-ACA derivative or by the decrease of ammonium consumption rate to 2 % of the initial consumption rate, the solution is filtered and the resin, in which glutaryl-7-ACA acylase is immobilised, is washed with, preferably 100 mM phosphate buffer pH 7.0.

10

The yield of 3-thiolated-7-ACA (compound II) after both enzymatic stages is about 94 %. The latter compound is then recovered from the final solution by crystallisation at an acidic pH from 1.7 to about 5.5, depending on the isoelectric point of the final product, with inorganic acids such as hydrochloric, sulphuric or phosphoric, or by combining any of these acids with suitable organic solvents at different ratios.

15

The process of the invention is new and permits a good overall product yield, better quality (in terms of colour and purity), easy isolation, a continuous operation of the process and the reuse of the enzymes avoiding the poisoning by the remaining heterocyclic thiol after chemical reaction.

20

25

The process according to the present invention represents a surprising improvement over known three-stage processes for obtaining 3-heterocyclic thiolated-7-aminocephalosporanic acid because cephalosporin C in solution is first reacted with a heterocyclic thiol.

This can be conveniently done on a solution of cephalosporin C derived directly from the fermentation broth. The process thus avoids the need to crystallise cephalosporin C as a metal salt, and, at the same time, eliminates the need for the use and recovery of organic solvents. Further, it reduces yield losses on the overall recovery process for cephalosporin C. Since the 3 substituted derivatives are more stable in solution than the parent cephalosporin C, overall fermentation yields are effectively increased still further.

The highly selective removal of excess thiol allows compounds of formula I to be prepared at a very high purity level with very low levels (<0.2 mg/ml) of heterocyclic thiols present. The process has several important advantages. It allows compounds of formula I to be used as a substrate for enzymatic processes without the poisoning of the enzymes. As a result the enzymes may be used repeatedly. In addition the process does not require the use of toxic reagents or the need to isolate intermediates thereby providing a continuous process.

A highly selective removal procedure with the strong anion exchanger Amberlite IRA-400 (manufactured by Rohm and Haas) is utilised, which allows the compounds of formula I with very low levels (<0.2 mg/ml) of heterocyclic thiols present to be obtained.

The reaction of nucleophilic substitution in the 3' position is carried out in an aqueous medium, dissolving the heterocyclic thiol and any non-toxic cephalosporin C salt in water by addition of a basic compound which form a water soluble salt, such as alkali metal hydroxide, ammonium hydroxide or preferably alkali metal carbonate or bicarbonate. In general, in addition to salts produced as described

above, any commercially available salt of cephalosporin C and of the heterocyclic thiols can be used in the process of this invention without changing the fundamentals of the process.

- 5 After dissolving the heterocyclic thiol and the cephalosporin C in separate reaction vessels or jointly, both reactants are mixed together in the same reactor, before or after heating the solution to a temperature from approximately 60°C to 80°C at a pH value of between 5.5 and 8.0.
- 10 Once the reaction starts, the temperature and pH are maintained preferably at approximately 65°C and 6.0 respectively, for a period of time of approximately 1 hour to 4 hours.
- 15 The heterocyclic thiol/cephalosporin C molar ratio is an important variable in the yield of the reaction and has to be optimised for each heterocyclic thiol used. Molar ratios are between 1.0 and 4.0, preferably at a molar ratio of approximately 4.
- 20 It was found that at these molar ratios the cephalosporin C remains quite stable with low β -lactam ring degradation, compared to a cephalosporin C solution without the thiol, which is completely degraded within 40 min at 80°C.
- 25 Once the cephalosporin C level is below 2% of the initial amount, the reaction mixture is cooled to a temperature from about 2°C to about 10°C, with or without acidification at a pH of from pH 3.0 to 5.5, preferably approximately 5.2, with strong mineral acids, such as hydrogen halides or oxy acids.

This acidification step gives in some cases, crystallisation of the heterocyclic thiol, with the concomitant possibility of reuse for a new reaction.

5

The resulting solution after the reaction with or without the acidifying step is subjected to subsequent purification by chromatography. Different resins and types of chromatography may be used on an industrial scale.

10

Several resins were tested grouped in four classes of resins based on adsorption, hydrophilic-hydrophobic interaction, cation exchange, and anion exchange. All resins tested based on adsorption (Amberlite XAD-761, Amberlite 7HP, Amberlite 16 HP and Amberlite XAD-4) gave similar results, the eluate containing from 22 % to 38 % of the heterocyclic thiol. The hydrophobic-hydrophilic interaction resin Sephadex LH-20 did not retain any thiol (<5%). A similar situation was found with the cation exchangers Amberlite ® IRC-50, IR-120 and IR-200. However, anion exchangers were found to have the best binding capacity for heterocyclic thiols ranging from 57-60 % in the case of a weak anion exchanger (Amberlite IRA-93).

15

20

25

It was found that a strong microporous (gel-type I) anion (base) exchange resin Amberlite IRA-400 having an 8% cross linking containing function trialkyl benzyl ammonium groups gave the highest binding of heterocyclic thiols (from 92-98 %) and low binding of the 3'-position heterocyclic thiomethyl cephalosporin C derivative (from 2-15 %, less than 15 % for the first cycle and less than 5 % for the following cycles).

Such a microporous resin offers certain advantages. They are less fragile, require less care in handling and possess higher loading capacities. As they have no discrete pores solute ions diffuse through the particle to interact with exchange sites. The total exchange capacity of the mentioned resin is in the order of 1,4 meq/ml.

It was surprisingly found that Amberlite IRA-400 has less binding capacity for 3-heterocyclic thiomethyl derivatives of cephalosporin C than for the same derivatives of glutaryl-7-ACA and 7-ACA. In fact the 3-heterocyclic thiomethyl derivative of glutaryl-7-ACA produced with MMTD binds at a level of 76.3 % to the column. The same result is found with 3-heterocyclic thiomethyl derivative of 7-ACA, with MMTD, which binds at a level of 92.7 % to the column. This unexpected behaviour of Amberlite IRA-400 with these three related β -lactam compounds appears to result from the presence of an ionisable amino group in the 5 position of the side chain of cephalosporin C compared with glutaryl-7-ACA and 7-ACA.

The removal of heterocyclic thiols by the process of the invention is particularly advantageous on an industrial scale as the elute of the column can be used for enzymation without isolation of the modified cephalosporin C and represents a new concept in the field of cephalosporin intermediates wherein the impurities are bound to the column and the β -lactam derivative is simply eluted by water.

Once the β -lactam derivative is eluted (less than 5 % remains bound), the column is typically regenerated with a 1.5 N solution of a strong mineral acid, such as hydrogen halide containing variable amounts of

an organic solvent, preferably 10-20 % acetonitrile. When the concentration of the thiol in the eluate is higher than 0.2 mg/ml, a strong regeneration using 3 N HCl and 40 % acetonitrile may be carried out. Alternatively, regeneration with successive NaOH and HCl solutions may be carried out.

After elution of the heterocyclic thiol, the thiol is concentrated and reused. The column is rinsed with deionised water to remove excess regenerant before the next cycle. The first bed volume of the rinse should be performed at the flow rate used for regeneration. The remainder is run at the adsorption flow rate.

The 3-thiolated derivative (TXC), has been found to be surprisingly good substrate, compared with unmodified cephalosporin C. The process of the present invention therefore provides an improved and more economical process for the preparation of cephalosporin C derivatives.

A second important aspect of the process of the invention is the use of enzymes in a re-usable form, either immobilised on a solid support or in the form of large stabilised crystals. This last requirement is important in producing a process to be operated on an industrial scale.

A further important advantage of the present invention is the ease of transferring the chemical solution after the removal and recovery of excess thiol to the first enzymatic reactor and from it to the second enzymatic reactor. This enables the process to be conducted continuously with a single liquid stream from cephalosporin C

concentrate or prepared batch solution to 3-thiolated-7-ACA derivative, which is easy to crystallise compared with unmodified 7-ACA.

5 The following examples are meant to illustrate the invention without limitation as to its generality.

Examples 1 to 5 illustrate the preparation of 3-thiolated-7-ACA derivatives of formula I from cephalosporin C.

10 Examples 6 to 10 illustrate the preparation of 3-thiolated-7-ACA derivatives of formula III through the formation of 3-thiolated-glutaryl-7-ACA derivatives of formula I.

15 **EXAMPLE 1:** Preparation of 7- β -(5-amino-5-carboxypentanamido)-3-(5-methyl-1,3,4-thiadiazole-2-yl thiomethyl)-3-cephem-4-carboxylic acid (TDC).

20 To a glass-lined reactor containing 600 ml of deionised water, 31.73 g (0.24 moles) of 2-mercapto-5-methyl-1,3,4-thiadiazole (MMTD) were added and the reactor was heated with stirring to a temperature of about 65°C. The pH of the mixture was adjusted to pH of about 6.0 by the addition of about 10 g of sodium carbonate.

25 In a separate glass-lined flask, a solution of concentrated sodium cephalosporin C (98 % purity by HPLC) was prepared by dissolving 33.23 g of sodium cephalosporin C (75 % free acid, 0.06 moles) in 200 mL of water. When the MMTD was dissolved, the concentrated cephalosporin C solution was added and the mixture was stirred at
30 approximately 65°C for 240 minutes, controlling the reaction kinetic

until the level of cephalosporin C was below 2 %. The following reaction kinetics were found:

Time (mins)	Cephalosporin C (moles)	TDC (moles)	MMTD (moles)
0	0.06	0.00	0.24
120	0.01	0.039	0.20
240	0.0012	0.042	0.195

5 The reaction mixture was then cooled to approximately 4°C, where the crystallisation of the excess MMTD begins. The pH was acidified with stirring (150 rpm) to a pH 5.2 with 37 % hydrochloride acid and left under slow stirring (50 rpm) for 60 minutes for the completion of crystallisation.

10

The precipitated MMTD was filtered and dried at 35 °C under vacuum. 23 g of recovered MMTD was obtained (purity 99 % by HPLC) with a recovery yield of about 95 %.

15

The filtrate (825 ml) containing 0.042 moles of the TDC and MMTD 0.016 moles was adjusted to pH 7.25 with 3 M ammonia and loaded onto an Amberlite IRA-400 column in chloride cycle (bed volume equal to 180 ml) covered with deionised water at flow rate 20 ml/min. Once loaded, the column was washed with deionised water (ca 100 ml) until 97 % recovery of loaded TDC with a 94 % purity by HPLC. The pH of the effluent was about 5.4 and was neutralised to 7.0 with 3 M ammonia. The remaining MMTD was 0.0009 moles (<0.2 mg/ml), which is less than 6 % of the remaining MMTD after its crystallisation by decreasing the pH. With this low level of MMTD (< 1 % of the

20

original MMTD after chemical reaction), enzymation of TDC is possible.

5 Typically the column is regenerated with 1 L of 1.5 M HCl containing 10 % acetonitrile and rinsed free of the excess regeneration by washing with 2 litres of deionised water. When required (MMTD > 0.2 mg/ml) the resin can be subjected to a strong regeneration using 1 litre of 3 M HCl with 40 % acetonitrile.

10 To further characterise the TDC solution at pH 5.0, it was loaded onto a Amberlite XAD-2 adsorption column and the column was washed with water. After washing, the resin was eluted with water, and 25 ml portions were pooled. A fraction containing 98.5 % TDC by HPLC was lyophilised and subjected to analysis:

15 Elemental Analysis for the product $C_{17}H_{20}N_5O_6S_3 \cdot 2H_2O$ (TDC), calculated C 37.42; H 4.43; N 12.84; S 17.63; found 37.27; H 4.3; N 13.11; S 17.51.

20 1H -NMR (DMSO- d_6) (δ ppm): 1.57 (m, 2H, $-CH_2-$); 1.63 (m, 2H, $-CH_2-$); 2.18 (m, 2H, $-CH_2-$); 2.64 (s, 3H, CH_3); 3.55, 3.73 (J=18 Hz, 2H, $-CH_2-$); 3.83 (t, 1H, $-CH-$); 4.18 – 4.46 (d, J=13 Hz, 2H, $-CH_2-$); 5.02 (d, J=3 Hz, 1H, C-6); 5.6 (d, J=3 Hz, 1H, C-7).

25

EXAMPLE 2: Comparative Example: Preparation of 7- β -(5-amino-5-carboxypentanamido)-3-(5-methyl-1,3,4-thiadiazole-2-yl thiomethyl)-3-cephem-4-carboxylic acid on different columns.

The TDC derivative was prepared as described in Example 1 and the filtrate containing it was loaded onto different types of resins.

5 The following data was obtained after washing with 100 ml of water in the first cycle of column usage:

Resin	Type	Eluted TDC (%)	Eluted MMTD (%)
Amberlite IRA-400	Strong anion exchanger	86	2
Diaion SA10A	Strong anion exchanger	67	15
Amberlite IRA93	Weak anion exchanger	87	45
Amberlite IRC-50	Weak cation exchanger	93	92
Amberlite IRC-200	Strong cation exchanger	73	98
Amberlite XAD-761	Adsorption	86	22
Amberlite XAD-7 HP	Adsorption	77	23
Amberlite XAD-16 HP	Adsorption	75	35
Amberlite XAD-4	Adsorption	68	25
Amberlite XAD-1180	Adsorption	78	38
Sephadex LH-20	Hydrophobic hydrophilic	98.5	95

Amberlite IRA-400 gave the best results. A high elution of TDC was observed with low elution of MMTD. Using other anion exchange columns higher

amounts of MMTD were also eluted. The other anion exchangers showed a high dual binding of thiol and TDC.

5 **EXAMPLE 3:** Specificity of TDC for Amberlite IRA-400.

The glutaryl-7-ACA derivative (TDG) and 7-ACA derivative (7-TDA) were prepared as in Example 1 using glutaryl-7-ACA and 7-ACA as starting material. The following data was obtained from the filtrate of the Amberlite IRA-400.

Resin	Compound in the elute	TDG or TDA eluted (%)	MMTD eluted (%)
Amberlite IRA-400	TDG	23.7	3.0
	TDA*	7.3	1.4

* At pH 7.8 to avoid any precipitation of TDA into the column.

15 In contrast to TDC, both TDG and TDA appear to remain bound to the Amberlite IRA-400 as well as the MMTD.

20 **EXAMPLE 4:** Preparation of 7- β -(5-amino-5-carboxypentanamido)-3-[(1-methyl-1H-tetrazol-5-yl)-thiomethyl]-cephalosporanic acid (TZC).

To a glass-lined reactor containing 600 ml of deionised water 28.16 g (0.24 moles) of 5-mercapto-1-methyltetrazole (MMTZ) were added and the reactor was heated with stirring to a temperature of approximately

70°C. The pH of the mixture was adjusted to pH of approximately 5.7-5.8 by the addition of approximately 12 g of sodium carbonate.

5 In a separate glass-lined flask, a solution of concentrated sodium cephalosporin C was prepared by dissolving 33.23 g of sodium cephalosporin C (75 % free acid, 0.06 moles purity 98 % by HPLC) in 200 ml of water. When the MMTZ was dissolved, the concentrated cephalosporin C solution was added and the mixture was stirred at about 70°C for 120 minutes, controlling the reaction kinetic until the
10 level of cephalosporin C was below 3 %.

Time (minutes)	Cephalosporin C (moles)	TZC (moles)	MMTZ (moles)
0	0.06	0	0.24
120	0.0017	0.040	0.19

15 The reaction mixture was cooled at about 4°C, but crystallisation of the excess of MMTZ did not start, even when the pH was decreased. The solution containing 0.04 moles of the TZC derivative from MMTZ and 0.19 moles of MMTZ was adjusted to pH 7.25 with 3 M ammonia and loaded onto an Amberlite IRA-400 column in chloride cycle (bed
20 volume equal to 150 ml) covered with deionised water at flow rate 20 ml/min. After the first pass through the column the remaining MMTZ was higher than 13 % of the initial (0.032 moles).

For this reason, the eluate was loaded onto another Amberlite IRA-400 (bed volume equal to 60 ml) column under the same conditions as described above to decrease the level of MMTZ.

Once loaded, the column was washed with deionised water (ca 90 ml) until 97 % recovery of loaded TZC with a 87% purity by HPLC. The pH of the effluent was about 5.4 and was neutralised to pH 7.0 with 3 M ammonia. The remaining MMTZ concentration was 0.0013 moles, which is less than 1 % of the original MMTZ after chemical reaction. With this low level of MMTZ, enzymation of the derivative is possible without poison the enzyme.

The columns were regenerated with 1 L of 1.5 M HCl containing 10 % acetonitrile and rinsed free of the excess regeneration by washing with 2 litres of deionised water.

EXAMPLE 5: Preparation of 7- β -(5-amino-5-carboxypentanamido)-3-[(1,2,5,6-tetrahydro-2-methyl-5,6-dioxo-1,2,4-triazin-3-yl)-thiomethyl]-cephalosporanic acid (TTC).

To a glass-lined reactor containing 600 mL of deionised water 37.96 g (0.24 moles) of 2,5-dihydro-3-mercapto-2-methyl-5,6-dioxo-1,2,4-triazine (here below indicated as TTZ) were added and the reactor was heated with stirring to a temperature of approximately 75°C. The pH of the mixture was adjusted to pH of about 6.7 by the addition of approximately 12 g of sodium carbonate.

In a separate glass-lined flask, a solution of concentrated sodium cephalosporin C was prepared by dissolving 33.23 g of sodium cephalosporin C (75 % free acid, 0.06 moles, purity 98 % by HPLC) in 200 ml of water. When the TTZ was dissolved, the concentrated

cephalosporin C solution was added and the mixture was stirred at about 75 °C for 75 minutes, controlling the reaction kinetic until the level of cephalosporin C was below 2 %.

Time (minutes)	Cephalosporin C (moles)	TTC (moles)	TTZ (moles)
0	0.06	0	0.24
75	0.0011	0.036	0.19

5 The reaction mixture was cooled at about 4°C, but crystallisation of the excess of TTZ does not start, even when the pH was decreased. The solution containing the 0.036 moles of TTC and 0.19 moles of TTZ was adjusted to pH 7.25 with 3 M ammonia and loaded onto an Amberlite IRA-400 column in chloride cycle (bed volume equal to 209 ml)
10 covered with deionised water at flow rate 20 ml/min. After the first column the remaining TTZ was 0.015 moles.

For this reason, the eluate was loaded onto another Amberlite IRA-400 column (with the same bed volume) under the same conditions as
15 described above to decrease the level of TTZ.

Once loaded, the column was washed with deionised water (ca 120 ml) until 60 % recovery of loaded TTC with a 90% purity by HPLC. The pH of the effluent was about 5.4 and was neutralised to pH 7.0 with 3 M
20 ammonia. The remaining TTZ concentration was 0.00096 moles, which is less than 1 % of the original TTZ after chemical reaction. With this level of TTZ, enzymation of the derivative is possible without poisoning the enzyme.

The columns were regenerated with 1 L of 1.5 M HCl containing 10% acetonitrile and rinsed free of the excess regeneration by washing with 2 litres of deionised water.

5

EXAMPLE 6: Synthesis of 7-amino-3-[(5-methyl-1,3,4-thiadiazol-2-yl)-thiomethyl]-cephalosporanic acid (TDA).

10 A filtrate (925 ml) from the strong anion exchanger Amberlite® IRA-400 from Example 1, containing 0.041 moles of 7 β -(5-amino-5-carboxypentamido)-3-[(5-methyl-1,3,4-thiadiazole-2-yl)-thiomethyl]-cephalosporanic acid, hereinbelow indicated as TDC, with 94.4 % purity (HPLC) and less than 0.2 mg/mL of 2-mercapto-5-methyl-1,3,4-
15 thiadiazole (MMTD) was adjusted to pH 7.0 with 3 M ammonia.

The TDC solution was fed into a 1.5 litre stirred reactor with 82.5 g (30.3 Roche's Units/g) of wet immobilised D-amino acid oxidase available from Roche Molecular Biochemicals and produced as
20 described in WO-A-9516773.

The conversion was performed at 20°C, 400 rpm and with an oxygen flow through a bottom diffuser of 0.1 vol/vol/min at 2 bar absolute pressure. The pH was titrated to pH 7.0 with 3 M ammonia by an
25 autotitrator.

The conversion was controlled by HPLC on a reverse phase column Nucleosil 120 3-C18 125 x 8 x 4 mm the mobile phase was 20 mM acetate ammonium pH 5.5 containing 4 % acetonitrile at 1 ml/min with

a 260 nm detector. The TDC appeared at 7.0 minutes, the α -ketoadipyl-intermediate at 8.5 min and the TDG (the 3-thiolated glutaryl-7-ACA) at 11.5 min.

- 5 Representative samples of the reaction mixture except for the enzyme were taken and the results obtained are reported in the following table:

	TDC	TDK	TDG	TDA	side products
Time (minutes)	[%] conversion	[%] production	[%] production	[%] production	[%] production
0	0.00	0.00	0.00	0.00	0.00
10	30.20	2.04	25.49	0.00	0.30
100	98.49	2.94	94.69	0.00	0.86

H ₂ O ₂ titration	98.60	0.00	97.71	0.00	0.89
---	-------	------	-------	------	------

- 10 When the conversion of TDC was higher than 98 %, the reaction was stopped and the reaction solution was filtered off. The remaining biocatalyst was washed with 100 ml of 100 mM phosphate buffer pH 7.0. This latter volume was added to the first filtrate.

- 15 To transform the residual α -ketoadipyl-thiolated derivative (TDK) to TDG, the solution obtained was titrated with 1 M hydrogen peroxide to 35 ppm for 30 minutes at 25°C. The conversion was controlled by HPLC until maximal production of TDG (see above table). The purity after treatment was: 92.65 % TDG and 7.34 % of other β -lactams.

The residual hydrogen peroxide was removed by adding 10 μ l of soluble *Corynebacterium glutamicum* catalase (650 kU, available from Roche Molecular Biochemicals) for 5 minutes.

5

The resulting TDG solution was adjusted to pH 8.0 with 3 M ammonia and transferred to a 1.5 L stirred reactor containing 57.08 g (87.6 Roche's Units/g) of wet immobilised glutaryl-7-ACA acylase (available from Roche Molecular Biochemicals). The conversion was performed at 20°C, 250 rpm and with a nitrogen flow through a bottom diffuser of 0.01 vol/vol/min at ambient pressure. The pH value was titrated to pH 8.0 with 3 M ammonia by an autotitrator. The conversion was controlled by HPLC with the same conditions as D-AAO reaction. The retention time for the product, 7-amino-3-[(5-methyl-1,3,4-thiadiazol-2-yl)-thiomethyl]-cephalosporanic acid (herebelow referred as TDA) was 7.2 minutes.

10

15

Representative samples of the reaction mixture were taken and the result obtained are shown in the following table:

20

	TDG	TDA	side products
Time (minutes)	[%] conversion	[%] production	[%] production
0	0.00	0.00	0.89
10	68.92	65.12	2.52
50	97.40	93.61	2.52

When the TDG conversion was higher than 97 %, the reaction was stopped and the reaction solution was filtered off. The remaining enzyme was washed with 100 ml of 100 mM phosphate buffer pH 7.0 and the remaining catalyst was ready for reuse. This latter volume was added to the filtrate, containing TDA, and cooled to 10°C. The pH was adjusted to 5.2 by concentrated sulphuric acid and allowed to crystallise for 60 minutes under slow stirring.

The precipitate was filtered off and washed successively with 250 ml of 50 % acetone/water and 100 ml of acetone, filtered again and dried at 35°C under vacuum to obtain 13.91 g (K.F. 1.81 %) of substantially pure 7-amino-3-[(methyl-1,3,4-thiadiazol-2-yl)-thiomethyl]-cephalosporanic acid, having an HPLC purity of 98.2 % and a transmittance at 420 nm of 87 (determined as 1 % solution in 2 % NaHCO₃; 1 cm cell).

The overall yield from the starting cephalosporin C solution (24.92 activity grams of free acid) in this CEE process to TDA was 64.6 %.

TDA

$^1\text{H-NMR}$ (DMSO/HCl) (δ ppm): 2.59 (s, 3H, $-\text{CH}_3$); 3.68 (broad s, 2H, $-\text{CH}_2-$); 4.2-4.48 ($J=13.2$ Hz, 2H, $-\text{CH}_2-$); 5.03 (d, $J = 4.8$ Hz, 1H, C-6); 5.11 (d, $J = 4.8$ Hz, 1H, C-7).

5 Elemental analysis:

Calculated for $\text{C}_{11}\text{H}_{12}\text{N}_4\text{S}_3\text{O}_3$: C:38.36 %; H: 3.51 %; S: 27.93 %; N: 16.27 %.

Found: C:38.35 %; H: 3.4 %; S: 28.00 %; N: 16.04 %.

10

EXAMPLE 7: Effect of heterocyclic thiol present in TDC solution in the reuses of the enzyme

15 To show the poisonous effect of remaining heterocyclic thiol in the two-step enzymation of TDC, TDC was used with and without heterocyclic thiol removal by chromatography in a strong anion exchanger Amberlite® IRA-400. The level of MMTD after crystallisation step was 2.56 mg/ml. After the chromatographic step onto an
20 Amberlite® IRA-400 strong anion exchanger, the amount of MMTD was less than 0.2 mg/mL, which represents less than 1% of the original MMTD used.

25 The following table shows how without the chromatographic step the length of the D-amino acid oxidase time doubles after 4 cycles, whereas it remains the same (≈ 100 minutes), after 12 cycles in the case of TDC with traces of MMTD.

Without chromatographic step		
CYCLE	D-amino acid oxidase	glutaryl-7-ACA acylase
	T(min)	T(min)
1	100	70
2	150	50
3	150	60
4	230	60
With chromatographic step		
CYCLE	D-amino acid oxidase	glutaryl-7-ACA acylase
	T(min)	T(min)
1	90	60
2	90	60
3	100	50
4	90	50
5	100	50
6	90	50
7	100	50
8	100	50
9	100	50
10	100	50
11	100	50
12	100	50

As regards, the second enzyme glutaryl-7-ACA-acylase, it seems that the hydrogen peroxide produced and added after the first reaction destroys traces of MMTD, preserving its catalytic activity with reaction times of about 50 minutes.

EXAMPLE 8: Synthesis of 7-amino-3-[(1-methyl-1H-tetrazol-5-yl)-thiomethyl]-cephalosporanic acid (TZA)

- 5 A filtrate (1000 ml) from the strong anion exchanger Amberlite ® IRA-400 from Example 4, containing 0.039 moles of 7-(5'-amidoadipamido)-3-[(1-methyl-1H-tetrazol-5-yl)-thiomethyl]-cephalosporanic acid (TZC) with 91.3 % purity (HPLC) and less than 0.2 mg/ml of 5-mercapto-1-methyltetrazole (MMTZ) was adjusted to pH 7.25 with 3 M ammonia.
- 10 The TZC solution was fed into a 1.5 litre stirred reactor with 82.5 g (30.3 Roche's Units/g) of wet immobilised D-amino acid oxidase available from Roche Molecular Biochemicals and produced as disclosed in WO-A-9516773.
- 15 The conversion was performed at 20°C, 400 rpm and with an oxygen flow through a bottom diffuser of 0.1 vol/vol/min at 2 bar absolute pressure. The pH was titrated to pH 7.25 with 3 M ammonia by an autotitrator.
- 20 The conversion was controlled by HPLC on a reverse phase column Nucleosil 120 3 - C18 125 x 8 x 4 mm; the mobile phase was 20 mM ammonium acetate pH 5.5 with 4% acetonitrile at 1 ml/min with a 260 nm detector. The TZC appeared at 3.0 minutes, the α -ketoadipyl-intermediate at 3.6 min and the TZG at 4.6 min (the 3-thiolated
- 25 glutaryl-7-ACA).

Representative samples of the reaction mixture except for the enzyme were taken and the result obtained are reported in the following table:

	TZC	TZK	TZG	TZA	side products
Time (minutes)	[%] conversion	[%] production	[%] production	[%] production	[%] production
0	0.00	0.00	0.00	0.00	0.00
10	73.98	6.49	66.02	0.00	0.30
35	100.00	4.26	91.28	0.00	2.12
H ₂ O ₂ titration	100.00	0.00	98.17	0.00	1.83

When the conversion of TZC was higher than 99 %, the reaction was stopped and the reaction solution was filtered off. The remaining biocatalyst was washed with 100 ml of 100 mM phosphate buffer pH 7.0. This latter volume was added to the first filtrate.

To transform the residual α -keto adipyl-thiolated derivate (TZK) to TZG, the solution obtained was titrated with 1 M hydrogen peroxide to 35 ppm for 30 minutes at 25°C. the conversion was controlled by HPLC (see above table). The purity after the treatment was: 89.45 % TZG and 10.55 % of other β -lactams.

The residual hydrogen peroxide was removed by adding 10 μ l of soluble *Corynebacterium glutamicum* catalase (650 kU, available from Roche Molecular Biochemicals) for 5 minutes.

The resulting TZG solution was adjusted to pH 8.0 with 3 M ammonia and transferred to a 1.5 L stirred reactor contained 57.08 g (87.6

Roche's Units/g) of wet immobilised glutaryl-7-ACA acylase (available from Roche Molecular Biochemicals). The conversion was performed at 20°C, 250 rpm and with a nitrogen flow through a bottom diffuser of 0.01 vol/vol/min at ambient pressure. The pH value was titrated to pH 8.0 with ammonia by an autotitrator. The conversion was controlled by HPLC with the same conditions as D-AAO reaction. The retention times for the product, 7-amino-3-[(5-methyl-1,3,4-thiadiazol-2-yl)-thiomethyl]-cephalosporanic acid (herein-below referred as TZA) was 3.2 minutes.

Representative samples of the reaction mixture were taken and the result obtained are shown in the following table

	TZK	TZG	TZA	side products
Time (minutes)	[%] conversion	[%] conversion	[%] production	[%] production
0	0.00	0.00	0.00	1.83
10	0.00	73.08	69.72	3.36
45	0.00	100.00	95.12	4.88

When the TZG conversion was higher than 99 %, the reaction was stopped and the reaction solution was filtered off. The remaining enzyme was washed with 100 ml of 100 mM phosphate buffer pH 7.0 and the remaining catalyst was ready for reuse. This latter volume was added to the filtrate, containing TZA, and cooled to 10°C. The pH was adjusted to 5.2 by concentrated sulphuric acid and allowed to crystallise for 60 minutes under slow stirring.

The precipitate was filtered off and washed successively with 250 mL of 50 % acetone/water and 100 ml of acetone, filtered again and dried at 35°C under vacuum to obtain 9.94 g (K. F. 1.4 %) of substantially pure 7-amino-3-[(methyl-1,3,4-thiadiazol-2-yl)-thiomethyl]-cephalosporanic acid, having a HPLC purity of 96.60 %.

The overall yield from the starting cephalosporin C solution (24.92 activity grams of free acid) in this ECC process to TZA was 48.05 %.

10 TZA

¹H-NMR (D₂O/Na₂CO₃) (δ ppm): 3.45-3.78 (J = 17.7 Hz, 2H, -CH₂-); 4.01-4.3 (J = 13.2 Hz, 2H, -CH₂-); 4.03 (s, 3H, -CH₃); 4.72 (d, J = 4.8 Hz, 1H, C-6); 5.00 (d, J = 4.8 Hz, 1H, C-7).

EXAMPLE 9: Effect of heterocyclic thiol present in TZC solution in the reuses of the enzyme

20 To show the poisonous effect of remaining heterocyclic thiol in the two-step enzymation of TZC, TZC was used with and without heterocyclic thiol removal by chromatography in a strong anion exchanger Amberlite® IRA-400. The level of MMTZ after chemical reaction step was 27.6 mg/ml After the chromatographic step onto an
25 Amberlite® IRA-400 strong anion exchanger was 0.2 mg/ml, which represent less than 1 % of MMTZ used.

The following table shows how without the two chromatographic steps, the length of the D-amino acid oxidase reaction increased a little

in two cycles, but dramatically affected the glutaryl-7-ACA acylase, which was unable to finish the second reaction in the second cycle.

Without chromatographic step		
CYCLE	D-amino acid oxidase	glutaryl-7-ACA acylase
	T(min)	T(min)
1	100	>110
2	120	>>110
With chromatographic step		
CYCLE	D-amino acid oxidase	glutaryl-7-ACA acylase
	T(min)	T(min)
1	45	45
2	35	45
3	35	40
4	40	48
5	45	48

However, the time of the cycles with chromatographic step, for both enzymes remained about the same in the five cycles presented in above table. This clearly shows the need of thiol removal for the stability of the enzymes.

EXAMPLE 10: Preparation of 7-amino-3-[(1,2,5,6-tetrahydro-2-methyl-5,6-dioxo-1,2,4-triazin-3-yl)-thiomethyl]-cephalosporanic acid (TTA).

A filtrate (1000 ml) from the strong anion exchanger Amberlite® IRA-400 from Example 5, containing 0.032 moles of 7 β -(5-amino-5-carboxypentamido)-3-[(1,2,5,6-tetrahydro-2-methyl-5,6-dioxo-1,2,4-triazin-3-yl)-thiomethyl]-cephalosporanic acid, hereinbelow indicated as TTC) with 88 % purity (HPLC) and less than 0.2 mg/ml of 2,5-

dihydro-3-mercapto-2-methyl-5,6-dioxo-1,2,4-triazine (TTZ) was adjusted to pH 7.25 with 3 M ammonia.

5 The TTC solution was fed into a 1,5 litre stirred reactor with 82.5 g (30.3 Roche's Units/g) of wet immobilised D-amino acid oxidase available from Roche Molecular Biochemicals and produced as disclosed in WO 95/16773.

10 The conversion was performed at 20°C, 400 rpm and with an oxygen flow through a bottom diffuser of 0.1 vol/vol/min at 2 bar absolute pressure. The pH was titrated to pH 7.25 with 3 M ammonia by an autotitrator.

15 The conversion was controlled by HPLC on a reverse phase column Nucleosil 100-5 C18 250 x 8 x 4.6 mm; the mobile phase was 25 % methanol in 10 mM tetrabutylammonium hydrogen sulfate and 15 mM potassium dihydrogen phosphate at 1 ml/min with a 260 nm detector. The TTC appeared at 4.3 minutes, the α -ketoadipyl-intermediated at 6.8 min and the 3-thiolated glutaryl-7-ACA (TTG) at 8.2 min.

20

Representative samples of the reaction mixture except for the enzyme were taken and the result obtained are reported in the following table:

	TTC	TTK	TTG	TTA	side products
Time (minutes)	[%] conversion	[%] production	[%] production	[%] production	[%] production
0	0.00	0.00	0.00	0.00	0.00
40	75.43	1.80	66.67	0.00	5.21

90	97.60	2.67	88.54	0.00	6.39
----	-------	------	-------	------	------

H ₂ O ₂ titration	97.60	0	90.96	0.00	6.5
--	-------	---	-------	------	-----

When the conversion of TTC was higher than 97 %, the reaction was stopped and the reaction solution was filtered off. The remaining biocatalyst was washed with 100 ml of 100 mM phosphate buffer pH 7.0. This latter volume was added to the first filtrate.

To transform the residual α -ketoadipyl-thiolated derivative (TTK) to TTG, the solution obtained was titrated with 1 M hydrogen peroxide to 35 ppm for 30 minutes at 25 °C. The conversion was controlled by HPLC (see above table). The purity after of treatment was: 80.77 % TTG and 19.23% of other β -lactams.

The residual hydrogen peroxide was removed by adding 10 μ l of soluble *Corynebacterium glutamicum* catalase (650 kU, available from Roche Molecular Biochemicals) for 5 minutes.

The resulting TTG solution was adjusted to pH 8.0 with 3 M ammonia and transferred to a 1,5 L stirred reactor contained 57.08 g (87.6 Roche's Units/g) of wet immobilised glutaryl-7-ACA acylase (available from Roche Molecular Biochemicals). The conversion was performed at 20°C, 250 rpm and with a nitrogen flow through a bottom diffuser of 0.01 vol/vol/min at ambient pressure. The pH value was titrated to pH 8.0 with ammonia by an autotitrator.

The conversion was controlled by HPLC with the same conditions as D-AAO reaction. The retention times for the product, 7-(5'-amidoadipamido)-3-[(1,2,5,6-tetrahydro-2-methyl-5,6-dioxo-1,2,4-triazin-3-yl)-thiomethyl]-cephalo-sporanic acid (TTA) was 5.2 minutes.

5

Representative samples of the reaction mixture were taken and the result obtained are shown in the following table:

	TTK	TTG	TTA	side products
Time (minutes)	[%] conversion	[%] conversion	[%] production	[%] production
0	0	0.00	0.00	6.39
10	0	94.33	84.49	7.13
50	0	100.00	89.09	7.64

When the TTG conversion was higher than 99 %, the reaction was stopped and the reaction solution was filtered off. The remaining enzyme was washed with 100 mL of phosphate buffer pH 7.0 and the remaining catalyst was ready for reuse. This latter volume was added to the filtrate, containing TTA, and cooled to 10 °C. The pH was adjusted to 4.2 by concentrated hydrochloric acid and allowed to crystallise for 60 minutes under slow stirring.

The precipitate was filtered off and washed successively with 250 ml of 50 % acetone/water and 100 ml of acetone, filtered again and dried at 40°C under vacuum to obtain 8.53 g (K.F. 5.95 %) of substantially pure 7-(5'-amidoadipamido)-3-[(1,2,5,6-tetrahydro-2-methyl-5,6-dioxo-1,2,4-triazin-3-yl)-thiomethyl]-cephalosporanic acid, having an HPLC purity of 98.28 %.

The overall yield from the starting cephalosporin C solution (24.92 activity grams of free acid) in this ECC process to TTA was 40.01 %.

TTA

^1H -NMR ($\text{D}_2\text{O}/\text{Na}_2\text{CO}_3$) (δ ppm): 3.43-3.7 ($J = 17.7$ Hz, 2H, $-\text{CH}_2-$); 3.63 (s, 3H, $-\text{CH}_3$); 4.02-4.33 ($J = 13.8$ Hz, 2H, $-\text{CH}_2-$); 4.73 (d, $J = 4.5$ Hz, 1H, C-6); 5.02 (d, $J = 4.5$ Hz, 1H, C-7).

- 5 The invention is not limited to the embodiments hereinbefore described which may be varied in detail.

10

15